

## Advanced glycation end products impair protein turnover in LLC-PK<sub>1</sub>: Amelioration by trypsin

GUANGSHENG XIANG, REINHARD SCHINZEL, ANDREAS SIMM, KATARINA SEBEKOVA, and AUGUST HEIDLAND

Department of Internal Medicine, Institute of Physiological Chemistry, and Institute of Clinical Biochemistry and Pathobiochemistry, University of Würzburg, Würzburg, Germany; and Institute of Preventive and Clinical Medicine, Bratislava, Slovakia

### Advanced glycation end products impair protein turnover in LLC-PK<sub>1</sub>: Amelioration by trypsin.

**Background.** Advanced glycation end products (AGEs) are assumed to play a key role in the pathogenesis of diabetic nephropathy (DN) and other diabetic complications. While AGEs have been shown to exert marked effects on mesangial and endothelial cells as well as on monocytes/macrophages, little is known about their effects on tubule cells. Therefore, we addressed the questions of (1) whether AGE-bovine serum albumin (AGE-BSA) impairs the protein metabolism in the tubule cells, and if so, (2) whether the AGE-induced effects are mediated via a protease sensitive mechanism.

**Methods.** Arrested LLC-PK<sub>1</sub> cells were exposed to a medium containing the vehicle (control, serum free), AGE-BSA (38  $\mu\text{mol/L}$ ), or BSA (38  $\mu\text{mol/L}$ ) in the presence or absence of trypsin (2.5  $\mu\text{g/mL}$ ) for 24 hours. We evaluated cell number, cell size, and cell protein content, as well as protein synthesis and protein degradation.

**Results.** After an incubation period of 24 hours, AGE-BSA decreased the cell number to  $84.5 \pm 5.5\%$  of control and  $82.5 \pm 5.6\%$  of BSA-treated cells ( $P < 0.05$ ). [<sup>3</sup>H]-thymidine incorporation declined to 66% of control ( $P < 0.05$ ), while BSA was without any effect. The same AGE-BSA dose reduced protein degradation ( $P < 0.05$ ) and stimulated total protein synthesis slightly, as determined by L-[<sup>14</sup>C]Phe incorporation into acid-insoluble proteins. These effects resulted in a rise in cell protein content (AGE-BSA vs. control,  $21.9 \pm 6.7\%$ ; AGE-BSA vs. BSA,  $11.1 \pm 6.0\%$ ,  $P < 0.05$ ) and cell volume (AGE-BSA vs. control  $9.4 \pm 3.2\%$ , AGE-BSA vs. BSA  $18.4 \pm 3.7\%$ ,  $P < 0.05$ ). Coincubation with AGE-BSA and trypsin was associated with an amelioration of all investigated parameters concerning cell number, cell proliferation, raised cell protein content, decreased protein degradation, and enhanced protein synthesis.

**Conclusion.** These data indicate that AGE-BSA impairs cell proliferation and protein turnover in LLC-PK<sub>1</sub> cells with a consequent rise in cell protein. Since these alterations were abrogated by coincubation with trypsin, an interference of this serine protease with the AGE-binding proteins on cell surfaces is assumed.

Diabetic nephropathy (DN) is characterized by renal hypertrophy, thickening of basement membranes, and accumulation of extracellular matrix (ECM) in the mesangium and tubulointerstitium [1, 2], and results both from enhanced production of proteins as well as from impaired protein degradation (PD) [3–6]. In the pathogenesis of DN, in addition to hyperglycemia, enhanced formation of transforming growth factor- $\beta_1$  (TGF- $\beta_1$ ) and activation of the renin-angiotensin system (RAS), early and advanced glycation end products (AGEs) are assumed to play a key role [7–11]. AGEs accumulate in the glomeruli and tubulointerstitium in relationship to severity of renal disease [12, 13], and administration of AGE-BSA to normal rats and mice led to lesions similar to those of DN [14, 15]. In *in vitro* studies, AGEs elicited a multitude of toxic effects such as enhanced formation of oxygen radicals, release of cytokines (in particular TGF- $\beta_1$ ), expression of adhesion molecules, and enhanced vascular permeability [16–18]. These responses most likely are mediated by AGE receptors, among which the receptor for AGEs (RAGE) is the best studied member [19–21]. Binding of AGEs to RAGE results in the activation of the transcription factor nuclear factor- $\kappa\text{B}$  as well as activator protein-1 (AP-1) complex [22–24].

In contrast to the numerous studies performed in mesangial and endothelial cells, there are only minimal data concerning AGE effects on tubular cells, the main target of renal hypertrophy in diabetes. Therefore, in the current study, the influence of AGE-BSA on proliferation and protein turnover of tubule cells was investigated. Furthermore, the effect of coadministered trypsin was examined, since AGE-binding proteins at surfaces of endothelial cells were shown to be sensitive to this serine protease [25, 26]. Thereby, an inactivation of the AGE receptor may protect the cells from maladaptive responses.

Results of the present study showed that exposure of LLC-PK<sub>1</sub> cells to AGE-BSA reduced cell number and

**Key words:** diabetic nephropathy, tubule cells, protein metabolism, cell proliferation, serine protease.

© 2001 by the International Society of Nephrology

[<sup>3</sup>H]-thymidine incorporation. Cell protein content and cell volume were increased because of impaired PD and slightly enhanced protein synthesis (PS). Coadministration of trypsin abrogated these alterations, most likely due to an interference of this serine protease with binding proteins of AGE-BSA at cell surfaces.

## METHODS

### Generation and characterization of AGE-modified albumin

Bovine serum albumin (1 mmol/L; BSA fraction V, Sigma, Germany) was glycosylated by incubation with 1 mol/L glucose in 50 mmol/L potassium phosphate, pH 7.3, 1 mmol/L ethylenediaminetetraacetic acid (EDTA) under sterile conditions at 50°C for 40 days according to a modified protocol of Schmidt et al [25].

### Cell culture and experimental treatment

LLC-PK<sub>1</sub> cells, a porcine cell line that exhibits properties of proximal tubules, were grown at 37°C in a humidified atmosphere of 5% CO<sub>2</sub> in Dulbecco's modified Eagles medium (DMEM) supplemented with 10% fetal calf serum (FCS; GIBCO, Grand Island, NY, USA), 25 mmol/L HEPES, 100 U/mL penicillin, and 100 ng/mL streptomycin. Cells were plated separately on 100 mm dishes (isolation of RNA), 6-well plates (cell proliferation assay), or 12-well plates (protein turnover measurements). Subconfluent cells were synchronized by serum-free medium (SFM) for 24 hours, followed by treatment with vehicle (control), BSA or AGE-BSA (19 to 76 μmol/L) related to BSA content with or without trypsin (0.625 to 5 μg/mL final concentration) for 24 hours. Cytotoxicity was evaluated both by trypan blue exclusion and release of lactate dehydrogenase (LDH). Cell number and cell volume was analyzed using the Casy-1 System (Schaefer, Reutlingen, Germany).

### [<sup>3</sup>H]-thymidine incorporation assay

The subconfluent monolayer was quiescented by incubation with serum-free DMEM for 24 hours and then incubated for another 24 hours in SFM in the absence (control) or the presence of AGE-BSA or nonglycosylated BSA with or without trypsin at various concentrations. During the last four hours, cells were labeled with 0.5 mCi/well of [<sup>3</sup>H]-thymidine (50 μCi-mmol; Amersham, Frankfurt/M, Germany). The cells were rapidly washed three times with ice-cold phosphate-buffered saline (PBS) and solubilized in 2% sodium dodecyl sulfate (SDS), followed by precipitation with 1 mL of 20% trichloroacetic acid. The precipitates were collected with a cell harvester onto a glass microfiber filter (Schleicher and Schuell, Dassel, Germany) and washed sequentially with 10 and 5% trichloroacetic acid (TCA) and finally

with ethanol. Incorporated [<sup>3</sup>H]-thymidine was measured by a liquid scintillation counter.

### Cellular protein content

Total protein concentration in cell lysate was measured using bicinchoninic acid (BCA protein assay; Pierce, Rockford, IL, USA) [27].

### Protein synthesis

Protein synthesis was assessed by determining the incorporation of L-[<sup>14</sup>C] phenylalanine (Phe; Amersham) into acid-insoluble proteins [28]. Subconfluent cells were preincubated for two hours in media containing 0.6 mmol/L unlabeled Phe, AGE-BSA, unmodified BSA, or vehicle (as control) before changing to a medium containing L-[<sup>14</sup>C] Phe (0.5 μCi/well) plus 0.6 mmol/L unlabeled Phe for 24 hours. The excess of unlabeled Phe (0.6 mmol/L) was used to ensure the equilibrium of intracellular- and extracellular-specific radioactivity. At the end of the four-hour labeling period, the cells were rapidly rinsed three times with ice-cold PBS. The cells were then solubilized in 2% SDS and precipitated. The radioactivity in the cells was determined in a Tri-Carb 4000 liquid scintillation counter (Packard Instruments, Downers Grove, CA, USA). The cell PS rate was expressed as dpm per 10<sup>5</sup> cells.

### Protein degradation

To measure PD rate, the quiescent subconfluent cells were labeled by 0.5 μCi/well of L-[<sup>14</sup>C]Phe for 24 hours [28]. To minimize the reincorporation of L-[<sup>14</sup>C]Phe, the experimental media contained an excess (2 mmol/L) of unlabeled Phe. After two hours, the cells were washed with ice-cold PBS to remove L-[<sup>14</sup>C]Phe released from the degradation of short-lived proteins; 4 mL of fresh experimental medium (control) containing BSA or AGE-BSA were added with and without trypsin. At the indicated time points, 0.5 mL aliquots of the media were collected and precipitated by adding ice-cold TCA (10% final concentration). The samples were centrifuged for five minutes, and the TCA-soluble supernatants were stored at -20°C. At the end of an experiment, the cells were washed three times with ice-cold PBS. Radioactivity in cells and media was measured as mentioned previously in this article. The percentage of radioactivity released versus the radioactivity remaining in the cells at the indicated time intervals was calculated as the PD rate and expressed as cpm/10<sup>5</sup> cells.

### Statistical analysis

The mean for the replicates of each experiment was determined. Results are presented as mean ± SD, with *N* indicating the number of experiments. A comparison between the two groups was made using paired and unpaired *t*-test. Group differences (>3) were evaluated em-

**Table 1.** Effect of different treatments on cell number, protein content and cell size after 24 hours of incubation

	Cell number $\times 10^6$	Protein content ng/cell	Cell volume $fL/10^3$
Control	$3.95 \pm 0.12$	$0.730 \pm 0.07$	$2827.79 \pm 101.6$
AGE-BSA	$3.34 \pm 0.22^a$	$0.894 \pm 0.06^a$	$3092.06 \pm 101.6^a$
BSA	$4.05 \pm 0.35$	$0.806 \pm 0.01$	$2611.85 \pm 96.7$
AGE-BSA + Tryp	$4.01 \pm 0.30$	$0.772 \pm 0.05$	$2632.97 \pm 97.56$
BSA + Tryp	$4.23 \pm 0.24$	$0.810 \pm 0.08$	$2865.89 \pm 110.1$

LLC-PK<sub>1</sub> cells were grown in vehicle, advanced glycation end products-bovine serum albumin (AGE-BSA) (38  $\mu$ mol/L), and BSA (38  $\mu$ mol/L) with or without trypsin (Tryp; 2.5  $\mu$ g/mL) for 24 hours. Cell monolayers were harvested by 0.25% trypsin/0.02% EDTA to obtain a single cell suspension, and subjected to the measurements of cell volume and cell number. Values are expressed as the mean  $\pm$  SD from three independent experiments ( $N = 9$ ).

<sup>a</sup> $P < 0.05$  vs control, BSA, AGE-BSA + Tryp

playing analysis of variance. Statistical significance was defined at  $P < 0.05$ .

## RESULTS

### Effect of AGE-BSA on cell proliferation

With AGE-BSA (38  $\mu$ mol/L), treatment of LLC-PK<sub>1</sub> cells, cell number was reduced within 24 hours to  $84.5 \pm 5.6\%$  in comparison to control ( $P < 0.05$ ,  $N = 9$ ) and to  $82.5 \pm 5.5\%$  in comparison to BSA-treated cells ( $P < 0.05$ ,  $N = 9$ ; Table 1). This reduction was not due to cell death, since both exclusion of trypan blue and release of LDH were not significantly changed in the different groups. [<sup>3</sup>H]-thymidine incorporation decreased within 24 hours from  $2678 \pm 170$  control to  $1857 \pm 178$  cmp/well ( $P < 0.05$ ). Coadministration of trypsin enhanced the incorporation to  $2321 \pm 214$  cmp/well ( $P < 0.05$ ).


### Effect of AGE-BSA on cell volume and cellular protein content

In the cells grown for 24 hours in AGE-BSA (38  $\mu$ mol/L), volume increased by  $9.4 \pm 3.2\%$  ( $P < 0.05$ ) as compared with untreated cells and by  $18.4 \pm 3.7\%$  ( $P < 0.05$ ) as compared with BSA-treated cells (Table 1). Coincubation of the cells with trypsin (2.5  $\mu$ g/mL) reversed this AGE-BSA effect.

Advanced glycation end product-BSA (38  $\mu$ mol/L) for 24 hours increased cellular protein content by  $21.9 \pm 6.7\%$  as compared with control ( $0.89 \pm 0.06$  vs.  $0.73 \pm 0.07$  ng/cell,  $P < 0.05$ ) and by  $11.1 \pm 6.0\%$  as compared with BSA-treated cells ( $0.89 \pm 0.06$  vs.  $0.80 \pm 0.01$  ng/cell,  $P < 0.05$ ). Protein content of cells grown in BSA was not significantly altered as compared with untreated cells. Coincubation of AGE-BSA with trypsin (2.5  $\mu$ g/mL) abolished the rise of cell protein content.

### Effect of AGE-BSA on PS

To exclude binding of AGE-BSA to cells as the main reason for the measured increase in cellular protein content, effects of AGE-BSA on cellular PS and degradation were analyzed in addition. [<sup>14</sup>C]-Phe incorporation into TCA-precipitated cell protein was slightly increased in



**Fig. 1.** Effect of advanced glycation end product-bovine serum albumin (AGE-BSA) and BSA on intracellular protein degradation in LLC-PK<sub>1</sub> cells. Exposure to AGE-BSA (38  $\mu$ mol/L) decreased protein degradation to 83% of controls after 24 hours. Coincubation of AGE-BSA with trypsin antagonized the AGE effect. Data are given as means  $\pm$  SD of eight determinations from two separate experiments. \* $P < 0.05$  vs. BSA-treated, AGE + trypsin-treated cells.

cells exposed to AGE-BSA (38  $\mu$ mol/L) in comparison to cells exposed to BSA (38  $\mu$ mol/L). The labeling incorporation rate after 12 and 24 hours was  $117.6 \pm 2.2\%$  ( $P < 0.05$ ) and  $109.9 \pm 10.5\%$  ( $P = NS$ ), respectively, when compared with the same concentration of BSA. Coincubation with trypsin (2.5  $\mu$ g/mL) prevented the rise in PS.

### Effect of AGE-BSA on protein degradation

After cells exposure to AGE-BSA (38  $\mu$ mol/L) for 24 hours, the release of incorporated [<sup>14</sup>C]-Phe was reduced by  $17 \pm 7\%$  as compared with the control ( $P < 0.05$ ,  $N = 6$ ) and by  $10 \pm 6\%$  as compared with BSA-treated cells ( $NS$ ,  $N = 6$ ; Fig. 1). Coincubation of the cells with trypsin (2.5  $\mu$ g/mL) increased the rate of PD by  $37 \pm 5.7\%$  as compared with AGE-BSA treatment alone ( $P < 0.05$ ,  $N = 6$ ).

## DISCUSSION

In the current investigation, AGE-BSA and not BSA inhibited the proliferation of LLC-PK<sub>1</sub> cells, as demonstrated by a decrease in cell number and [<sup>3</sup>H]-thymidine incorporation without any effects on cell viability and LDH release. These findings are consistent with studies in mesangial [29, 30], smooth muscle [31], and endothelial cells [32], as well as retinal capillary pericytes [32].

The antiproliferative action of AGE-BSA in LLC-PK<sub>1</sub> cells was associated with a rise in cell volume and cell protein content. PS (as demonstrated by incorporation of L-<sup>14</sup>C Phe) was slightly increased, while PD was significantly reduced. The latter finding might be explained, at least in part, by impaired proteolytic activities. Thus, in our earlier study, incubation of LLC-PK<sub>1</sub> cells with AGE-BSA caused a marked decline of cathepsin B + L activities caused by lowered enzyme expression [33].

These alterations are most probably caused by an AGE-BSA-induced overexpression of TGF- $\beta_1$ , which was shown to enhance PS on the one hand and to impair PD on the other [34, 35]. Indeed, in our own preliminary studies, TGF- $\beta_1$  overexpression was observed in porcine LLC-PK<sub>1</sub> cells after AGE-BSA incubation (abstract; Xiang et al, *J Am Soc Nephrol* 9:629, 1998), corresponding with data in mesangial cells [16]. Additionally, in vivo administration of AGE-BSA in normal rats and mice led to an overexpression of TGF- $\beta_1$  in the glomeruli [15].

To prevent the intracellular/extracellular effects of AGEs, various experimental and clinical approaches have been employed. Administration of aminoguanidine, a blocker of AGE formation, attenuated the renal pathology of streptozotocin (STZ)-diabetic rats [36]. Additionally, antioxidants (taurine, vitamin E, and alpha-lipoic acid) protected mesangial and/or endothelial cells from AGEs and/or glucose induced injury, probably by decreased activation of nuclear factor- $\kappa$ B and/or TGF- $\beta_1$ , respectively [24, 37]. Recently, administration of the soluble receptor of AGEs was reported to ameliorate diabetic atherosclerosis [38].

In our study, we investigated the question of whether trypsin modulates the effect of AGE-BSA in LLC-PK<sub>1</sub> cells, since this protease inhibits the protein binding of AGEs at the surface of endothelial cells [25, 26]. Our data show that coincubation of AGE-BSA with trypsin ameliorates the antiproliferative effects, the raised cell protein content, as well as the increased PS, and impairs PD. In preliminary investigations, we could also observe a decrease of intracellular AGE accumulation in LLC-PK<sub>1</sub> cells, as determined by an immunohistochemical staining method of imidazolone, which could explain the beneficial effects of trypsin on the investigated parameters (data not shown).

The question arises, how could AGEs exert their toxic effects in tubular cells in DN? Contact with the basolat-

eral membrane is determined by the blood concentration of AGEs and at the apical site by the glomerular filtration of the low molecular weight AGE peptides. Thus, after injections of AGE containing peptides in normal rats, a reabsorption of these compounds by proximal tubules has been demonstrated [39]. Moreover, in the presence of glomerular proteinuria, besides AGE peptides the higher molecular weight AGE proteins also are translocated into the proximal tubular fluid. Here, they may interact with cell receptors at the apical site with consequent cell activation. Recently, binding sites specific for AGEs were demonstrated within the proximal tubule cells of the rat kidney [40].

In conclusion, our data show that AGE-BSA exerts marked effects on LLC-PK<sub>1</sub> cells, such as decreased proliferation, rise in cell volume and cell protein content, impaired PD, and slightly enhanced PS. These alterations are substantially inhibited by coadministration of trypsin. It is assumed that this serine protease hinders the interaction of AGEs with cell binding proteins, thereby preventing the release of cytokines.

Reprint requests to August Heidland, M.D., Department of Internal Medicine, University of Würzburg, Josef-Schneider-Strasse 2, 97080 Würzburg, Germany.

## REFERENCES

1. ZIYADEH FN: The extracellular matrix in diabetic nephropathy. *Am J Kidney Dis* 22:736-744, 1993
2. MAUER SM: Structural-functional correlations of diabetic nephropathy. *Kidney Int* 45:612-618, 1994
3. RECKELHOFF JF, TYGART VL, MITIAS MM, WALCOTT JL: Decreased activity of glomerular cathepsin and metalloproteinases in the rat. *Diabetes* 42:1425-1432, 1993
4. SCHAEFER L, SCHAEFER RM, LING H, TESCHNER M, HEIDLAND A: Renal proteinases and kidney hypertrophy in experimental diabetes. *Diabetologia* 37:567-571, 1994
5. SHECHTER P, BONER G, RABKIN R: Tubular cell protein degradation in early diabetic renal hypertrophy. *J Am Soc Nephrol* 4:1582-1587, 1994
6. PACZEK L, SOIN J, BARTOLOMIEJCZYK I, KROLAK R, LAO M, GRADOWSKA L, HEIDLAND A: Impairment of proteolytic activity in isolated glomeruli and tubules in experimental diabetic nephropathy. *Clin Nephrol* 46:258-260, 1996
7. YAMAMOTO T, NAKAMURA T, NOBLE NA, RUOSLAHTI E, BORDER WA: Expression of transforming growth factor beta is elevated in human and experimental diabetic nephropathy. *Proc Natl Acad Sci USA* 90:1814-1818, 1993
8. SHARMA K, JIN Y, GUO J, ZIYADEH FN: Neutralization of TGF- $\beta$  by anti-TGF- $\beta$  antibody attenuates kidney hypertrophy and the enhanced extracellular matrix gene expression in STZ-induced diabetic mice. *Diabetes* 45:522-530, 1996
9. WOLF G, SHARMA K, CHEN Y, ERICKSEN M, ZIYADEH FN: High glucose induced proliferation in mesangial cells is reversed by autocrine TGF- $\beta$ . *Kidney Int* 42:647-656, 1992
10. COHEN MP, ZIYADEH FN: Role of amadori-modified nonenzymatically glycosylated serum proteins in pathogenesis of diabetic nephropathy. *J Am Soc Nephrol* 7:183-190, 1996
11. BROWNLEE M, CERAMI A, VLISSARA H: Advanced glycosylation end products in tissue and the biochemical basis of diabetic complications. *N Engl J Med* 318:1315-1321, 1988
12. BEISSWENGER PJ, MAKITA Z, CURPHEY TJ, MOORE LL, JEAN S, BRINCK-JOHNSON T, BUCALA R, VLISSARA H: Formation of immunohistochemical advanced glycosylation end products precedes and cor-

- relates with early manifestations of renal and retinal disease in diabetes. *Diabetes* 44:824-829, 1995
13. BENDAYAN M: Immunocytochemical detection of advanced glycation end products in rat renal tissue as a function of age and diabetes. *Kidney Int* 54:438-447, 1998
  14. VLASSARA H, STRIKER LJ, TEICHBERG S, FUH H, LI YM, STEFFES M: Advanced glycosylation end products induce glomerular sclerosis and albuminuria in normal rats. *Proc Natl Acad Sci USA* 91:11704-11708, 1994
  15. YANG CW, VLASSARA H, PETEN EP, HE CJ, STRIKER GE, STRIKER LJ: Advanced glycation end products up-regulate gene expression found in diabetic glomerular disease. *Proc Natl Acad Sci USA* 91:9436-9440, 1994
  16. THROCKMORTON DC, BROGDEN AP, MIN B, RASMUSSEN H, KASHGARIAN M: PDGF and TGF-beta mediate collagen production by mesangial cells exposed to advanced glycosylation end products. *Kidney Int* 48:111-117, 1995
  17. BAYNES JW, THORPE SR: Perspectives in diabetes: Role of oxidative stress in diabetic complications: A new perspective on an old paradigm. *Diabetes* 48:1-9, 1999
  18. MIYATA T, IIDA Y, UEDA Y, SHINZATO T, SEO H, MONNIER VM, MAEDA K, WADA K: Monocyte/macrophage response to beta 2-microglobulin modified with advanced glycation end products. *Kidney Int* 49:538-550, 1996
  19. YAN SD, SCHMIDT AM, ANDERSON GM, ZHANG J, BRETT J, ZOU YS, PINSKY D, STERN D: Enhanced cellular oxidant stress by the interaction of advanced glycation end products with their receptor-binding proteins. *J Biol Chem* 269:9989-9997, 1994
  20. NEEPER M, SCHMIDT AM, BRETT J, YAN SD, WANG F, PAN YC, ELLISTON K, STERN D, SHAW A: Cloning and expression of a cell surface receptor for advanced glycosylation end products of proteins. *J Biol Chem* 267:14998-15004, 1992
  21. GRETEN J, KREIS I, WIESEL K, STIER E, SCHMIDT AM, STERN DM, RITZ E, WALDHERR R, NAWROTH PP: Receptors for advanced glycation end products (AGE)-expression by endothelial cells in nondiabetic uraemic patients. *Nephrol Dial Transplant* 11:796-802, 1996
  22. LANDER HM, TAURAS JM, OGISTE JS, HORI O, MOSS RA, SCHMIDT AM: Activation of the receptor for advanced glycation end products triggers a p21-ras-dependent mitogen-activated protein kinase pathway regulated by oxidant stress. *J Biol Chem* 272:17810-17814, 1997
  23. SIMM A, MÜNCH G, SEIF F, SCHENK O, HEIDLAND A, RICHTER H, VAMVAKAS S, SCHINZEL R: Advanced glycation stimulate the MAP-kinase pathway in tubules cell line LLC-PK1. *FEBS Lett* 410:481-484, 1997
  24. BIERHAUS A, CHEVION S, CHEVION M, QUEHENBERGER P, HOFMANN M, ILLMER T, LUTHER T, BERENTSSTEIN E, TRITSCHLER H, MÜLLER M, WAHL P, ZIEGLER R, NAWROTH PP: Advanced glycation end products (AGEs) induced activation of NF- $\kappa$ B is suppressed by  $\alpha$ -lipoic acid in cultured endothelial cells. *Diabetes* 46:1481-1490, 1997
  25. SCHMIDT AM, VIANNA M, GERLACH M, BRETT J, RYAN J, KAO J, STERN D: Isolation and characterization of two binding proteins for advanced glycation end products from bovine lung which are present on the endothelial cell surface. *J Biol Chem* 267:14987-14997, 1994
  26. YAN SD, CHEN Y, FU J, CHEN M, SCHMIDT AM: RAGE and amyloid-beta peptide neurotoxicity in Alzheimer's disease. *Nature* 382:685-691, 1996
  27. SMITH PK, KROHN RJ, HERMANSON GT, MALLIA AK, GARTER FH, PROVENZANO MD, FUJIMOTO EK, GOEKE NM, OLSON BJ, KLENK DC: Measurement of protein using bicinchoninic acid. *Anal Biochem* 150:76-85, 1985
  28. JURKOVITZ CT, ENGLAND BK, EBB RG, MITCH WE: Influence of ammonia and pH on protein and amino acid metabolism in LLC-PK<sub>1</sub> cells. *Kidney Int* 42:595-601, 1992
  29. CROWLEY STM, BROWNLEE M, EDELSTEIN JA, SATRIANO T, MORI P, SINGHAL P, SCHLÖNDORFF D: Effects of nonenzymatic glycosylation of mesangial matrix on proliferation of mesangial cells. *Diabetes* 40:540-547, 1991
  30. IEHARA N, TAKEOKA H, YOSHIHIKO Y, KITA T, DOI T: Advanced glycation end products modulate transcriptional regulation in mesangial cells. *Kidney Int* 50:1166-1172, 1996
  31. IINO K, YOSHINARI M, YAMAMOTO M, KAKU K, DOI Y, ICHIKAWA K, IWASE M, FUJISHIMA M: Effect of glycated collagen on proliferation of human smooth muscle cells in vitro. *Diabetologia* 39:800-806, 1996
  32. CHIBBER R, MOLINATTI PA, ROSATTO N, LAMBOURNE B, KOHNER EM: Toxic action of advanced glycation end-products on cultured retinal capillary pericytes and endothelial cells: Relevance to diabetic retinopathy. *Diabetologia* 40:156-164, 1997
  33. SEBEKOVA K, SCHINZEL R, LING H, SIMM A, XIANG G, GEKLE M, MÜNCH G, VAMVAKAS S, HEIDLAND A: Advanced glycation albumin impairs protein degradation in the kidney proximal tubules cell line LLC-PK<sub>1</sub>. *Cell Mol Biol* 44:1051-1060, 1998
  34. BORDER WA, NOBLE NA: Transforming growth factor- $\beta_1$  in tissue fibrosis. *N Engl J Med* 331:1286-1292, 1994
  35. LING H, VAMVAKAS S, BUSCH G, DÄMMRICH J, SCHRAMM L, LANG F, HEIDLAND A: Suppressing role of transforming growth factor- $\beta_1$  on cathepsin activity in cultured tubule cells. *Am J Physiol* 269(6 Pt 2):F911-F917, 1995
  36. SOULIS-LIPAROTA T, COOPER M, PAPAZOGLU D, CLARKE B, JERUMS G: Retardation by aminoguanidine of development of albuminuria, mesangial expansion and tissue fluorescence in streptozotocin-induced diabetic rat. *Diabetes* 40:1328-1334, 1991
  37. TRACHTMAN H, FUTTERWEIT S, PRENNER J, HANON S: Antioxidants reverse the antiproliferative effect of high glucose and advanced glycosylation end products in cultured rat mesangial cells. *Biochem Biophys Res Commun* 199:346-352, 1994
  38. PARK L, RAMAN KG, LEE KJ, LU Y, FERRAN LJ Jr, CHOW WW, STERN D, SCHMIDT AM: Suppression of accelerated diabetic atherosclerosis by the soluble receptor for advanced glycation end products. *Nat Med* 4:1025-1031, 1998
  39. GUGLIUCCI A, BENDAYAN M: Renal fate of circulating advanced glycation end products (AGE): Evidence for reabsorption and catabolism of AGE-peptides by renal proximal tubular cells. *Diabetologia* 39:149-160, 1996
  40. YOUSSEF S, NGUYEN DT, SOULIS T, PANAGIOTOPOULOS S, JERUMS G, COOPER ME: Effect of diabetes and aminoguanidine therapy on renal advanced glycation end-product binding. *Kidney Int* 55: 907-916, 1999